The ageing kidney/ Chronic kidney failure

Nele Van Den Noortgate IUC – 17th May 2018

Content

- Ageing of the kidney
 - Structural and functional changes
 - Mechanisms of age-dependent injury
- Chronic Kidney Disease in the older person
 - Methods to estimate GFR
 - Is a decline in GFR equivalent to CKD?
 - Managing CKD in the older person



Structural changes in the ageing kidney

- Macroscopy
 - Decrease weight
 - Decrease size
 - Cortical atrophy

- Microscopy
 - Glomerular
 - Tubular
 - Interstitial
 - Vascular

Progressive loss of renal mass

- Loss of cortical mass due to fall in total number of glomeruli
- Tubular and interstitial changes
- Vascular changes

Age specific incidence of glomerulosclerosis





Kaplan et al , Am J Pathol 80:227-234,1975

Glomerular changes

- Thickening of the basement membrane
- Reduplication of Bowman's capsule
- Mesangial matrix expansion
- Fusion of foot processes



Decline in GFR

Baltimore Longitudinal study

Age (years)	No subjects	creat clearance (ml/min/1.73m ²)	creat clearance slope (ml/min/1.73m²/yr)
45-54	95	128.1±1.6	-0.73 ± 0.30
55-64	60	121.8±1.9	-1.64 ± 0.41
65-74	36	110.0±2.6	-1.30 ± 0.57
75-84	17	97.0±3.4	-1.07 ± 0.77

Rowe et al. J Gerontol 1976;31:155-163

Tubular and interstitial changes

- Reduction in proximal tubular volume and number
- Irregular thickening of basal membrane
- Interstitial changes consists of increasing zone of tubular atrophy and fibrosis



Electrolyte disorders in the old old in an acute geriatric ward (220 pts)

- hypokalaemia (K < 3.5 mmol/l): 26 patients
- hyperkalaemia (K > 5.5 mmol/l): 9 patients
- hyponatraemia (Na < 135 mmol/l): 46 patients
- hypernatraemia (Na > 145 mmol/l): 4 patients
- significant correlation between hypokalaemia and locomotoric disturbances (p= 0.01); the use of diuretics(p=0.005), laxatives (p=0.006); failure to thrive (p=0.005)

Van Den Noortgate et al. Int Urol Nephrol 2001;32:531-537.

Effect of abrupt salt restriction



Epstein et al. J Lab Clin Med 1976;87:411-17

Effect of salt-loading



Epstein et al. J Lab Clin Med 1976;87:411-17

The concentrating capacity of the kidney

Urine osmolality in response to 12 hoursof water deprivation

Cyclic AMP concentrations in renal papillary slices after vasopressin



Rowe et al. Nephron 1976;17:270-78



Serum K and aldosterone levels before, during and after infusion with KCI

Mulkerrin et al JASN 6: 1459-1462, 1995

Vascular changes

- Fibrous intima thickening
- Loss of media
- Arteriolar hyalinosis



Content

- Ageing of the kidney
 - Structural and functional changes
 - Mechanisms of age-dependent injury
- Chronic Kidney Disease in the older person
 - Methods to estimate GFR
 - Is a decline in GFR equivalent to CKD?
 - Managing CKD in the older person



Etiology of ageing (kidney)

Interactions between

- genetically individual susceptibility
- intrinsic stresses
- extrinsic factors

High heterogeneity (in GFR) in the elderly

Heterogeneity in GFR in the elderly





Etiology of the ageing kidney

- Genetic events
 - modulation of lifespan by a large number of genes
 - replicative senescence is determined by shortening of telomeres, cell cycle inhibitory genes, senescence associated beta-galactosidase
 - markers of senescence also present in the ageing kidney

Regression of telomeric length in renal cortex in relation to age



Melk et al JASN, 11: 444-453, 2000



Westhoff et al. JASN 2010;21:327-36

Etiology of the ageing kidney

- Environmental and lifestyle factors
 - Diseases influencing GFR
 - hypertension
 - vascular disease
 - diabetes mellitus
 - heart failure

Frohling et al. Kidney Int 1989;36(suppl 27): 91-95

Etiology of the ageing kidney



Renal vascular resistance in the aged kidney

Increased sympathetic tone Stimulated norepinephrine levels

Decrease in prostaglandins

Autoregulatory defence in the aged kidney

Increased Intrarenal vascular tone Angiotensin II ↑ Endogenous endothelin ↑ Nitric Oxide ↓

Content

- Ageing of the kidney
 - Structural and functional changes
 - Mechanisms of age-dependent injury
- Chronic Kidney Disease in the older person
 - Methods to estimate GFR
 - Is a decline in GFR equivalent to CKD?
 - Managing CKD in the older person



Aline - 85 year old

- Medical history
 - Arterial hypertension
 - Ischemic heart disease CABG in 2010
- Current history
 - Fall at home with subcapital humerus fracture
 - Living independently
 - Only cleaning aid



- Medication
 - Asaflow 80 mg
 - Simvastatine 40 mg
 - Perindopril 5 mg
 - Co-Bisoprolol 5/12,5 mg
- Medical examen
 - G: 50 kg; L: 1m65; BMI: 18
- Biochemistry
 - Creatinine 1,0 mg/dl (2j terug idem)

Does Aline suffer from decline in eGFR?



Methods to measure renal function

• Renal function is estimated by GFR, measured as the renal clearance of a particular substance:

GFR = UV/P

- Substances used
 - Exogenous (gold standard) impractical & expensive
 - ⁵¹Cr-EDTA; inulin
 - Endogenous
 - Serumcreatinine and cystatin C

Methods to estimate GFR: serum creatinine

- serum creatinine/ reciprocal serum creatinine
 - muscle mass declines with ageing
 - assay for serum creatinine (compensated Jaffé)
- urinary creatinine clearance
 - urine collections
 - tubular secretion of creatinine
- mathematical corrections (age, gender, weight)
 - Cockcroft-Gault formula
 - MDRD-formula
 - CKD-EPI

Methods to measure GFR: Serum cystatin C

- non-glycosated, low molecular weight protein
- characteristics of an ideal GFR marker
 - endogenously produced at a constant rate
 - freely filtered at the glomerulus, destroyed in the proximal renal tubule, not extrarenally eliminated
 - not dependent on muscle mass, BMI, gender

Newman et al. Kidney Int 1995; 47:312-8.

e-GFR formulas in older persons

	Schattingsformules	voor renale klaring: mathematisch
CG	creatinineklaring =	(140 – leeftijd). Gewicht 72. Screat. (0,85 indien een vrouw)
MDRD	$GFR = 186.(Screat)^{-1.7}$	¹⁵⁴ . (leeftijd) ^{-0,203} . (0,742 indien vrouw) . (1,210 indien zwarte huidskleur)
CKD-EPI	$Vrouw + Screat \leq 0,7mg/dL$	$GFR = 144. (Screat/0,9)^{-0.329}. 0,993^{leeftijd}$
(blanke ras)	Vrouw + Screat > 0,7mg/dL	$GFR = 144.(Screat/0,9)^{-1,209}.0,993^{leeftijd}$
	$Man + Screat \leq 0,9mg/dL$	$GFR = 141.(Screat/0,9)^{-0,411}.0,993^{leeftijd}$
	Man + Screat > 0.9mg/dL	$GFR = 141.(Screat/0,9)^{-1,209}.0,993^{leeftijd}$
BIS	GFR	e = 3736. Screat ^{-0,87} . leeftijd ^{-0,95}

e-GFR formulas in older persons

- Cross-sectional analysis ; Geriatric ward UH
- 89 patients (25 men, 64 women)
 - Median age 85 years (80 89 years)
- renal evaluation on day 7 of admission
 - ⁵¹Cr-EDTA clearance
 - blood sample (serum creatinine-compensated jaffé, serum cystatin C, serum albumine, serum urea level)
 - 24h urine collections (measured creatinine clearance)
 - calculated clearance (Cockcroft-Gault & MDRD abbreviated)
- Second data-analyse for CKD-EPI&BIS in 2015

N Van Den Noortgate et al. JAGS 2002;50:1278-1282

Methdology BIAS: mean/median difference between eGFR - mGFR PRECISION: standaarddeviation (SD) of interquartielrange (IQR) of bias ACCURACY: a. % correct classified KDIGO-stadia b.% eGFR within 30% of the mGFR



Accuracy

Results

	total	CG	MDRD	CKD EPI	BIS
bias	mean	-3,0678	15,4112	7,2378	-0,2616
	median	-0,6528	14,1113	8,7611	4,8973
ision	IQR	[-9,78 / 6,73]	[2,50 / 29,74]	[-3,42 / 21,50]	[-11,04 / 11,46]
preci	SD	16,91	23,94	22,15	20,95
racy	P30	68,5%	40,4%	47,2%	66,3%
accur	stad	56/86 = 65,1%	38/89 = 42,7%	39/89 = 43,8%	45/89 = 50,5%

< 6	0 ml/min	CG	MDRD	CKD EPI	BIS
as	mean	1,9796	19,7118	14,0748	7,3599
pi	median	2,0569	19,0642	12,4710	7,0562
precisio	IQR	[-5,52 / 7,88]	[7,47 / 30,84]	[3,2 / 24,92]	[1,92 / 13,71]
	SD	9,89169	18,77556	15,74605	11,45014
racy	P30	75,4%	31,3%	34,3%	64,2%
accu	stad	47/67 = 70,1%	23/67 = 34,3%	24/67 = 35,8%	39/67 = 58,2%
≥ 6	0 ml/min	CG	MDRD	CKD EPI	BIS
as Se	0 ml/min mean	CG -18,6904	MDRD 2,3188	CKD EPI -13,5838	BIS -23,4725
bias ⊽	0 ml/min mean median	CG -18,6904 -19,6118	MDRD 2,3188 -0,2056	CKD EPI -13,5838 -9,3460	BIS -23,4725 -18,9491
cisio bias <	0 ml/min mean median IQR	CG -18,6904 -19,6118 [-32,5 / -7,03]	MDRD 2,3188 -0,2056 [-5,7 / 22,17]	CKD EPI -13,5838 -9,3460 [-22,47 / 2,14]	BIS -23,4725 -18,9491 [-32,84 / -11,3]
precisio bias <	0 ml/min mean median IQR SD	CG -18,6904 -19,6118 [-32,5 / -7,03] 23,73	MDRD 2,3188 -0,2056 [-5,7 / 22,17] 32,47530	CKD EPI -13,5838 -9,3460 [-22,47 / 2,14] 25,96735	BIS -23,4725 -18,9491 [-32,84 / -11,3] 26,03923
racy precisio bias ⊽	0 ml/min mean median IQR SD P30	CG -18,6904 -19,6118 [-32,5 / -7,03] 23,73 57,1%	MDRD 2,3188 -0,2056 [-5,7 / 22,17] 32,47530 68,2%	CKD EPI -13,5838 -9,3460 [-22,47 / 2,14] 25,96735 86,4%	BIS -23,4725 -18,9491 [-32,84 / -11,3] 26,03923 72,7%

Methods to evaluate renal function in elderly patients: a systematic literature review



Figure 1. Mean difference between the gold standard and the formula to calculate the GFR (with 95% CIs) for the separate studies. Top: GFR calculated with the CG formula. Bottom: GFR calculated with the MDRD formula.

Van Pottelberghe et al. Age Ageing 2010:39:542-548



Figure 2. Mean difference between the gold standard and the formula to calculate the GFR (with 95% CIs) for the separate studies. Top: Sensitivity of the CG and MDRD formulas, serum creatinine and cystatin C concentrations and creatinine clearance for the pooled data with cut-off values of 50, 60, 80 and 90 ml/min. Middle: Positive predictive value of the CG and MDRD formulas, serum creatinine and cystatin C concentrations and creatinine clearance for the pooled data with cut-off values of 50, 60, 80 and 90 ml/min. Bottom: Sensitivity of the CG and MDRD formulas, serum creatinine and cystatin C concentrations and creatinine clearance for the pooled data with cut-off values of 50, 60, 80 and 90 ml/min. Bottom: Sensitivity of the CG and MDRD formulas, serum creatinine and cystatin C concentrations and creatinine clearance for the cut-off value of 60 ml/min for pooled data and for the separate studies. CG, Cockcroft–Gault formula; MDRD, Modification of Diet in Renal Disease formula; Creat S, serum creatinine; Creat CL, creatinine clearance; cyst C, cystatin C.
Use of e-GFR methods in older persons

- Serum creatinine is an insensitive measure
- CG and BIS are of comparable value in an older hospitalized population with decline in GFR
- MDRD en CKD-epi are more useful to measure early decline in GFR
- Cystatin C is comparable with serum creatinine; it probably performs better in detecting very early decline in GFR

Content

- Ageing of the kidney
 - Structural and functional changes
 - Mechanisms of age-dependent injury
- Chronic Kidney Disease in the older person
 - Methods to estimate GFR
 - Is a decline in GFR equivalent to CKD?
 - Managing CKD in the older person



Impairment in eGFR with Ageing

Cross-sectional studies: NHANES III

GFR	Age Group (years)				
$mL/min/1.73m^2$ =	20 - 39	40 - 59	60 - 69	≥70	
≥90	86.0%	55.7%	38.5%	25.5%	
60 - 89	13.7%	42.7%	53.8%	48.5%	
30 - 59	*	1.8%	7.1%	24.6%	
15 - 29	*	*	*	1.3%	

* fewer than 20 cases; data not considered reliable

National Kidney Foundation. Am J Kidney Dis 2002; 39 (suppl 1): S58

Prevalence of reduced renal function- acute geriatric ward (UH-Ghent)

GFR estimated by MDRD	GFR estimated by CC-Gault
8.7	0
29.0	3.1
56.5	70.4
5.8	26.5
	<i>GFR estimated</i> <i>by MDRD</i> 8.7 29.0 56.5 5.8

Van Den Noortgate et al. Int Urol Nephrol 2001;32:531-537.

Does Aline suffer from Chronic Kidney Disease?



Table 1	Table 1. Stages of CKD ^a				
Stage	Description	GFR (mL/min/1.73 m²)			
1	Kidney damage with normal or GFR	≥ 90			
2	Kidney damage with mild GFR	89-60			
ЗА	Mild to moderate GFR	59-45			
ЗB	Moderate GFR	45-30			
4	Severe GFR	30-15			
5	Kidney failure	< 15 or dialysis			

CKD, chronic kidney disease; GFR, glomerular filtration rate.

*Adapted from the Renal Association. http://www.renal.org/whatwedo/InformationResources/ CKDeGUIDE/CKDstages.aspx. Accessed November 16, 2013.

Is decline in eGFR = normal ageing or CKD?

- No good correlation between RR, cardiac function and eGFR in older people
 - = do we measure normal ageing?

 Presence of proteinuria is an important indicator for CV risk and mortality in the older person

Risk of end-stage renal disease/ mortality in the older population with CKD?



El-Ghoul, Joly et al. JAGS 2009;57:2217-23

Diagnosis of CKD

eGFR < 60 ml/min

&

protein/creatinin ratio > 450 mg/g albumin/creatinin ratio (diabetes) > 30 mg/g

= kidney disease with high CV risk; not associated with normal ageing

Diagnosis of CKD

			Persist	ent albuminuria cobego Rescription and range)	éns	
				Al	A2	Að
Prognosis of CRD by BFR and albuminuria caleportex: KB468 2912		Ronmal to mildly increased	Modecately increased	Sevenets increased		
				< 30 mp/g < 3 mp/mmol	30-300 mg/g 300 mg/mmol	> 380 mg/g > 38 mg/ternol
	41	Normal or high	≥10			
Ê.	42	Wildly decreased	40-40			
Training of the local sector of the local sect	631	trially to moderately decrement	45-58			
topper la	608	Wederstelp to serverely decreased	30-04			
04.00	64	Severely decreased	15-25			
	65	Kidney failure	< 28			

Aline 85 years old

- Aline has an estimated GFR of
 - CG 32,5 ml/min
 - MDRD 53 ml/min
 - CKD-epi 51 ml/min
 Protein/creatinin ratio of 500 mg/g
- Management in this women?



Etiology of CKD



Nephrol Dial Transplant (2017) 32:9-16 doi: 10.1093/ndt/gfw411



NDT Perspectives

Clinical Practice Guideline on management of older patients with chronic kidney disease stage 3b or higher (eGFR<45 mL/min/1.73 m²): a summary document from the European Renal Best Practice Group

Ken Farrington¹, Adrian Covic², Ionut Nistor², Filippo Aucella³, Naomi Clyne⁴, Leen De Vos⁵, Andrew Findlay¹, Denis Fouque⁶, Tomasz Grodzicki⁷, Osasuyi Iyasere⁸, Kitty J. Jager⁹, Hanneke Joosten¹⁰, Juan Florencio Macias¹¹, Andrew Mooney¹², Evi Nagler⁵, Dorothea Nitsch¹³, Maarten Taal¹⁴, James Tattersall¹², Marijke Stryckers⁵, Dieneke van Asselt¹⁵, Nele Van den Noortgate¹⁶, Sabine van der Veer¹⁷ and Wim van Biesen⁵

- Treat possible etiologies of CKD
 - Diagnose/treat diabetes mellitus
 - Treat arterial hypertension SRR>140 mmHg?
 - Ultrasound of kidney
- Detect metabolic abnormalities
 - Hemoglobin/ 25 OH vit D
 - For people with eGFR < 30 ml/min
 - Bicarbonate/Calcium/phosphate/potassium
 - Parathormone

- Prevention of decline in kidney function
 - Start/continue nephroprotective medication
 - ACE-I in people with proteinuria
 - Avoid nephrotoxic medication/contrast
 - Adapt dose of medication with renal clearance (Cockroft-Gault)
 - Start diet low in salt
 - Stop smoking
 - Protein low diet?

• Detect frailty



Reese et al. Am J Nephrol. 2013;38:307-315

Mortality and Hospitalization	Intermediately Frail	Frail
Hazard ratio of mortality (95% CI)		
Unadjusted	2.67 (1.06-6.73)	2.90 (1.18-7.11)
Adjusted for age, sex, comorbidity, and disability	2.68 (1.02-7.07)	2.60 (1.04-6.49)
Incident rate ratio of hospitalization	(95% CI)	
Unadjusted	0.74 (0.49-1.12)	1.48 (1.05-2.07)
Adjusted for age, sex, comorbidity, and disability	0.76 (0.49–1.16)	1.43 (1.00-2.03)

Table 2. Mortality and Hospitalization According to Frailty Status (Reference Nonfrail)

CI = confidence interval.

Mc Adams-DeMarco et al. JAGS 2013;61:896-601 Johansen et al. JASN 2007;18:2960-2967

Riskfactors of falls

Table 2 Factors associated with falls in adults undergoing hemodialysis

	Unadjusted	Parsimonious model	Full model
Frailty status			12
Nonfrail	Reference	Reference	Reference
Intermediately frail	1.11 (0.44, 2.82)	1.35 (0.52, 3.49)	1.19 (0.44, 3.24)
Frail	3.55 (1.68, 7.46)**	3.89 (1.78, 8.49)**	3.09 (1.38, 6.90)**
Age (in 10 years)	-	0.83 (0.68, 1.01)	0.90 (0.69, 1.17)
Female	-5	1.68 (1.01, 2.79)*	1.84 (1.06, 3.19)*
Caucasian		0.98 (0.46, 2.12)	0.84 (0.37, 1.91)
Comorbidity ¹	8	8	0.88 (0.47, 1.65)
Disability ²	8	8	1.71 (0.97, 3.00)
Medication use ³	-	-	0.91 (0.85, 0.98)*
High school education or higher	2	2	3.63 (1.29, 10.24)*

Mara McAdams-DeMarco et al. BMC nephrology 2013;14:224

Q4A: WHAT IS THE BEST ALTERNATIVE METHOD TO ASSESS FUNCTIONAL DECLINE IN OLDER AND/OR FRAIL PATIENTS WITH ADVANCED CKD?

4a.1 We recommend a simple score be used on a regular basis to assess functional status in older patients with CKD stage 3b–5d with the intention to identify those who would benefit from more in-depth geriatric assessment and rehabilitation (1C)

4a.2 We recommend that most simple scores, including self-report scales and field tests (sit to stand, gait speed or 6-min walk test) have comparable and sufficient discriminating power to identify patients with decreased functional status (1C)

EB interventions in CKD

Q4B: ARE INTERVENTIONS AIMED AT INCREASING FUNCTIONAL STATUS IN OLDER PATIENTS WITH RENAL FAILURE (eGFR <45 ML/MIN/1.73 M² OR ON DIALYSIS) OF BENEFIT?

4b.1 We recommend that exercise has a positive impact on the functional status of older patients with CKD stage 3b or higher (1C) 4b.2 We suggest that exercise training be offered in a structured and individualized manner to avoid adverse events (2C)

5b.1 We suggest a trial of structured dietary advice and support with the aim of improving nutritional status (2C)

Casus Aline



- Normal metabolic parameters
- Frail and high risk for falls
- Refer to nephrologist?

Indications for referral to nephrologist

- eGFR lower than 30 ml/min
- eGFR between 30 45 ml/min and progressive decline in eGFR (> 5 ml/min/2j) or important proteinuria
 - Slower progression of kidney failure in older people:
 - 27% of 75y and older with baseline eGFR 25-55 ml/min has no decline in GFR over 10y
- Sudden decline in eGFR

Domus medica 2013 Rosansky et al. Am J Nephrol 2012;36:1-10

Indications for referral to nephrologist

Figure 2. Referral decision making by estimated glomerular filtration rate (eGFR) and albuminuria

				U	rine ACR categori scription and rar	les lige
				A1.	A2	A3 Severely increased
				Normal to mildly increased	Moderately increased	
-				<3mg/mmol	3-30mg/mmol	>30mg/mmol
- 14	G1	Normal or high	290		Monitor	Refer*
bue	63	Mildly decreased	60-89	()	Monitor	Refer*
tes (inter	G3a	Mildly to moderately decreased	45-59	Monitor	Monitor	Beter
(1.7.) /1.7.1 lona	G3b	Moderately to severely decreased	30-64	Monitor	Mannaty	Onlaw .
R al	64	Severely decreased	15-29	Refer*	Bullet .	Retim
218	65	Kidney failure	×15	in failer in		Malas
the second secon				the second se		

Frans 82 years old

- Medical history
 - Arterial hypertension
 - Diabetes mellitus since 10 years
 - Peripheral arterial disease
- Functioning
 - low activity (most of the time in chair/bed); help for washing; social isolated
- Kidney function
 - eGFR from 30 ml/min to 20 ml/min (over two years)
 - Albumin/creatinine ratio 30 mg/g
 - Hypovitaminose D en hyperparathyroidie



Frans 82 years old

- Dyspnoea and oedema
- Blood results
 - eGFR 10 ml/min
 - Bicarbonate 18 ml/min
 - Hyperparathyreose en hypovitaminose D
 - Hb 10,8 mg/dl
 - Phosphor slightly increased

Indication for RRT?



Q6: WHAT IS THE BENEFIT OF DIALYSIS IN FRAIL AND OLDER PATIENTS?

6.1 We recommend use of validated tools as explained in Questions 2 and 3 to project likely outcomes and help decide the appropriateness of discussing options for RRT (see Figure 1)6.2 We recommend that the option for conservative management be discussed during the shared decision-making process on different management options for ESKD (1D)

6.3 We recommend the REIN score can be useful to stratify short term/6 month mortality risk of patients intending to start RRT (1C)



FIGURE 1: Proposed management pathway for older patients with advanced CKD. KRFE score is the 4-variable Kidney Failure Risk Equation (see Question 2). For Bansal and REIN score see Question 3.

BANS score

Table 2

Parameter estimates in the final models and risk score

Variables	Parameter estimate in the final model	Ratio	Score
Older	0.96	2.97	3
DM	0.85	2.63	3
Hypotension	0.41	1.25	1
High CTR	0.38	1.16	1
High BNP level	0.32	1	1
ECG score = 1	0.46	1.43	1
ECG score = 2	1.52	4.69	5

Each parameter estimate in the final models was compared with the smallest parameter estimate (High BNP level). Then, the risk scores were determined.

Older 65 ≤ age, *DM* diabetes mellitus as a cause of end-stage renal disease, *Hypotension* hypotension during hemodialysis, *High CTR* 50 % ≤ cardiothoracic ratio, *High BNP level* 250 pg/mL ≤ plasma brain natriuretic peptide level, *ECG score* the number of abnormal findings in electrocardiogram

BANS SCORE

Survival probability



KRFE SCORE

Estimate risk of progression to end-stage renal disease in CKD patients using age, sex, eGFR and proteinuria with KFRE

Sex? Male Female Age? eGFR? Urine Albumin Creatinine Ratio? (Note units carefully) Patient location? North America Non-North America

REIN score

CG Couchoud et al.: Stratification of incident elderly ESRD patients

clinical investigation

	Mean a imputed	among 20 d data sets	Among 2000 samples (100 resampling)	Multivariate model	Score
Risk factors	N	% Death	Number of significant associations (P<0.05)	Adjusted odds ratio (95% confidence interval)	Points
Gender			1.000		
Male	7549	10.8	1944	1.23 (1.08-1.40)	1
Female	4951	9.7	-		
Age (years)					
(75-80)	5103	10.2	-	1	
(80-85)	4549	13.1	229	1.10 (0.95-1.27)	0
(85-90)	2393	18.1	2000	1.40 (1.19-1.66)	2
≥ 90		10.1	2000	1.79 (1.35–2.38)	3
Diabetes					
No	7437	10.1			
Yes	5063	10.8	2		
Congestive heart failure					
No	792	7.8	-	1	
Stage I-II	3185	12.9	2000	1.43 (1.23-1.67)	2
Stage III-IV	1395	19.7	2000	2.15 (1.79-2.59)	4
Peripheral vascular disease					
No or stage I-II	11.520	9.7	-		
Stage III-IV	980	18.2	1536	1.34 (1.09-1.64)	1
schemic heart disease					
No	8226	9.7	—		
Without myocardial infaction	2627	11.0	19		
With myocardial infaction	1648	13.1	213		
Cerebral vascular disease					
No	10.833	10.1	_		
Yes	1667	12.7	8		
199 State 1		100.00			

Table 2 Adjusted odds ratios for 3-month mortality in the training set, after imputation, and points assigned to each risk factor

Dysrhythmia					
No	8296	8.9	_	1	
Yes	4204	13.5	1939	1.25 (1.09–1.43)	1
Chronic respiratory disease					
No	10,624	10.0			
Yes	1876	12.8	2		
Cancer					
No	10,919	9.7	_	1	
Yes	1581	15.4	2000	1.61 (1.36–1.91)	2
Cirrhosis					
No	12,374	10.3	_		
Yes	126	18.3	326		
Severe behavioral disorder					
No	11,997	9.9	_	1	
Yes	503	21.4	1841	1.44 (1.12–1.85)	2
Mobility					
Walks without help	8502	5.8	_	1	
Need assistance for transfer	3159	15.9	2000	2.47 (2.10-2.91)	4
Totally dependent for transfer	839	36.6	2000	6.53 (5.38-7.92)	9
Albuminemia (g/l)					
<25	1175	21.4	2000	5.17 (2.94-2.22)	5
(25-30)	2468	14.4	2000	3.35 (2.01-1.66)	3
(30-35)	3987	10.2	1999	2.16 (1.57-1.30)	2
≥ 35	3379	6.4	_	1	
Body mass index (kg/m ²)					
<18.5	430	13.2	57		
(18.5-23)	2616	10.5	5		
(23-35)	1754	9.6	_		
(25-30)	3080	10.1	177		
≥ 30	1430	10.7	706		
Area under the curve				0.76 (0.75-0.77)	0.76 (0.75-0.77)



Figure 1 Distribution of the patients in the validation set, by score, according to vital status and dialysis withdrawal within 3 months.



Figure 2 Proposed risk stratification algorithm to screen for evaluation, evaluate and decide on the appropriate strategy of care for elderly ESRD patients, according to their level of risk of early death. CKD, chronic kidney disease.

Prognostic factors for RRT?

- Bad prognostic factors
 - Fast decline eGFR: 4 à 5 ml/min/year
 - Important proteinuria
- Most studies: 80-plus with CKD stage 4 died before indication to start RRT

Demoulin et al. NDT 2011;26:220-26 O hare et al. JASN 2007;18:2758-65

RRT and mortality in older people

Conservative Reference Dialysis group Results management group Carson et al., 200921 median age 83.0 median age 75.0 median survival from first known date of 13.8% diabetes 29.5% diabetes GFR $\leq 10.8^{a}$: mean age-adjusted CCI mean age adjusted CCI 13.9 months (range 2-44) with CM 37.8 months (range 0-106) with dialysis score 7.4 score 7.2 n = 29n = 173p < 0.01Chanda et al., 201017 mean age 77.5 mean age 58.5 median survival from first known date of 68.4% over age 75 11.2% over age 75 GFR < 15: 35.5% diabetes 34.3% diabetes 21.2 months with CMb 67.1 months with dialysis^b 49.7% high comorbidity 17.3% high comorbidity n = 155n = 689p < 0.001Ellam et al., 200919 median age 80 None median survival from first known date of 38% diabetes GFR < 15: 21 months (range 1-100) with CM 32% ischemic heart disease n = 69Joly et al., 2003²⁰ mean age 84.1 median survival from first day of dialysis mean age 83.2 51.4% late referral to 28.9% late referral to or decision not to perform dialysis: nephrology nephrology 8.9 months (95% CI 4-10) with CM 21.6% diabetes 6.5% diabetes 28.9 months (95% CI 24-38) with dialysis 48.6% ischemic heart disease 42.5% ischemic heart p < 0.000143.3% socially isolated disease n = 3714.7% socially isolated n = 107Murtagh et al., 200718 median age 79.6 median age 83 median survival from first known date of 23.4% diabetes 25.0% diabetes GFR < 15: n = 77n = 5218.0 months (range 0.1-73.1) with CM 19.6 months (range 2.2-84.2) with dialysis Smith et al., 2003¹⁵ $n = 34^{\circ}$ $n = 10^{\circ}$ median survival from proposed date of dialysis initiation: 6.3 months (range 0-46) with CM 8.3 months (range 2-20) with dialysis Wong et al., 200716 median age 79 median survival from decision not to None mean GFR 12 perform dialysis: 23.4 months with CMb 28% diabetes 34% ischemic heart disease n = 73

TABLE 2. PROGNOSIS WITH CONSERVATIVE, NONDIALYTIC MANAGEMENT OF END-STAGE RENAL DISEASE

O'Connor et al. J Pall Med 2012; 15:228-235

No benefit of RRT on mortality

Murtagh et al., 2007 ¹⁸	median age 83 23.4% diabetes	median age 79.6 25.0% diabetes	median survival from first known date of GFR < 15:
	n=77	<i>n</i> =52	18.0 months (range 0.1–73.1) with CM 19.6 months (range 2.2–84.2) with dialysis
Smith et al., 2003 ¹⁵	$n=34^{\circ}$	$n = 10^{\circ}$	median survival from proposed date of dialysis initiation:
			6.3 months (range 0–46) with CM 8.3 months (range 2–20) with dialysis

O'Connor et al. J Pall Med 2012; 15:228-235
Benefit of RRT on mortality

Reference	Conservative management group	Dialysis group	Results
Carson et al., 2009 ²¹	median age 83.0 13.8% diabetes mean age-adjusted CCI score 7.4 n=29	median age 75.0 29.5% diabetes mean age adjusted CCT score 7.2 n=173	median survival from first known date of GFR ≤10.8ª: 13.9 months (range 2–44) with CM 37.8 months (range 0–106) with dialysis p<0.01
Chanda et al., 2010 ¹⁷	mean age 77.5 68.4% over age 75 35.5% diabetes 49.7% high comorbidity n=155	mean age 58.5 11.2% over age 75 34.3% diabetes 17.3% high comorbidity n=689	median survival from first known date of GFR < 15: 21.2 months with CM ^b 67.1 months with dialysis ^b 9<0.001
Ellam et al., 2009 ¹⁹	median age 80 38% diabetes 32% ischemic heart disease n=69	None	median survival from first known date of GFR < 15: 21 months (range 1–100) with CM
Joly et al., 2003 ²⁰	mean age 84.1 51.4% late referral to nephrology 21.6% diabetes 48.6% ischemic heart disease 43.3% socially isolated n=37	mean age 83.2 28.9% late referral to nephrology 6.5% diabetes 42.5% ischemic heart disease 14.7% socially isolated n=107	median survival from first day of dialysis or decision not to perform dialysis: 8.9 months (95% CI 4–10) with CM 28.9 months (95% CI 24–38) with dialysis p < 0.0001

Benefit of RRT on mortality

Reference	Conservative management group	Dialysis group	Results
Carson et al.	, 2009 ²¹ median age 83.0	median age 75.0	median survival from first known date of
Joly et al., 2003 ²⁰	mean age 84.1 51.4% late referral to nephrology 21.6% diabetes 48.6% ischemic heart disease 43.3% socially isolated n=37	<pre>mean age 83.2 28.9% late referral to nephrology 6.5% diabetes 42.5% ischemic heart disease 14.7% socially isolated n=107</pre>	median survival from first day of dialysis or decision not to perform dialysis: 8.9 months (95% CI 4–10) with CM 28.9 months (95% CI 24–38) with dialysis p<0.0001
	51.4% late referral to nephrology 21.6% diabetes 48.6% ischemic heart disease 43.3% socially isolated n=37	 28.9% late referral to nephrology 6.5% diabetes 42.5% ischemic heart disease 14.7% socially isolated n=107 	or decision not to perform dialysis: 8.9 months (95% CI 4–10) with CM 28.9 months (95% CI 24–38) with dialysis p < 0.0001

Mortality in 80-plus: Risk factor



Figure 2. Actuarial survival curves in the group of hemodialyzed patients according to characteristics at inclusion. (a) Effect of gender on survival. (b) Survival in three groups defined by Karnofsky performance scale. (c) Effect of late referral. (e) Effect of peripheral vascular disease. (f) Effect of coronary artery disease. (g) Effect of cardiac failure.

Joly et al. J Am Soc Nephrol 2003;14: 1012-21

anarap) or concorridatio managomena

Aims: We aimed to compare survival, hospital admissions and palliative care access of patients aged over 70 years with chronic kidney disease stage 5 according to whether they chose renal replacement therapy or conservative management. **Design:** Retrospective observational study.

Setting/participants: Patients aged over 70 years attending pre-dialysis clinic.

Results: In total, 172 patients chose conservative management and 269 chose renal replacement therapy. The renal replacement







Distribution of Days Survived: Hospital-free Days, Outpatient Hemodialysis Days and Hospital Inpatient Days



Figure 3. Median survival for MCM cohort and the hemodialysis-only subgroup in the RRT cohort. Data shown are how many days were spent hospital-free, compared with in-patient stays in hospital and outpatient hospital attendances for dialysis.

Carson et al. Clin J Am Soc Nephrol 2009;4:1611-19

Kwaliteit van leven

Nephrology

Am J Nephrol 2013;37:231–238 DOI: <u>10.1159/000347220</u>

Received: December 1 2, 2012 Accepted: January 20, 2013 Published online: March 2, 2013

Trajectory of Quality of Life for Poor Prognosis Stage 5D Chronic Kidney Disease with and without Dialysis

Ying-Ying Seow^a Yin Bun Cheung^b Li Min Qu^c Alethea Chung Pheng Yee^c ^aKhoo Teck Phuat Hospital, ^bDuke-NUS Graduate Medical School, and ^cNational Cancer Centre, Singapore

Prospectieve studie – 101 patiënten eGFR 8 – 12 ml/min ≥ 75 jaar oud Charlson comorbidity index ≥ 8



Seow et al. JASN 2013;27:231-238

Renal replacement therapy

- Better outcome in
 - Highly active older people
 - Low comorbidity
 - Younger than 80 years
- Rather conservative in frail older people
 - Comprehensive geriatric evaluation
 - Multidisciplinary evaluation (GP/nephrologist/geriatrician)
 - Evaluation of needs/wishes of patient/family

Frans 82 years

- Started HD for his wife
- No quality hypotensive delirium fall
- Withdrawal of dialysis
- Died six month later at home



Take Home Message

- Renal Senescence is characterized by structural and physiological changes leading to reduced reserve capacity
- eGFR equations (CG, MDRD and CKD-EPI) are at this moment the best tools available to estimate renal function, however clinicians should be aware of their pitfalls
- Reduced eGFR should be distinguished from CKD
- CKD should be managed EB with prevention of decline of renal function, physical and nutritional support, regular evaluation through BANS & KRFE & REIN score
- Starting RRT should be a multidisciplinary decision together with patient and family. Conservative treatment can be in individual cases a good alternative.

